

Refine Search

Search Results -

Terms	Documents
L9 and night adj sight	1

Database:

US Pre-Grant Publication Full-Text Database
 US Patents Full-Text Database
 US OCR Full-Text Database
 EPO Abstracts Database
 JPO Abstracts Database
 Derwent World Patents Index
 IBM Technical Disclosure Bulletins

Search:

L11

Search History

DATE: Monday, September 27, 2004 [Printable Copy](#) [Create Case](#)

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=OR</i>			
<u>L11</u>	L9 and night adj sight	1	<u>L11</u>
<u>L10</u>	L9 and glare	5	<u>L10</u>
<u>L9</u>	L8 and @py>1975<=2002	55	<u>L9</u>
<u>L8</u>	L7 and eye	73	<u>L8</u>
<u>L7</u>	aceclidine	129	<u>L7</u>
<u>L6</u>	L5 and @py>1975<=2003	33	<u>L6</u>
<u>L5</u>	L4 and azelastine	43	<u>L5</u>
<u>L4</u>	antazoline	481	<u>L4</u>
<u>L3</u>	L2 and ophthalmic	38	<u>L3</u>
<u>L2</u>	L1 and pheniramine	98	<u>L2</u>
<u>L1</u>	ketotifen	960	<u>L1</u>

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptaul25zxf

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 Jul 12 BEILSTEIN enhanced with new display and select options,
resulting in a closer connection to BABS
NEWS 4 Jul 30 BEILSTEIN on STN workshop to be held August 24 in conjunction
with the 228th ACS National Meeting
NEWS 5 AUG 02 IFIPAT/IFIUDB/IFICDB reloaded with new search and display
fields
NEWS 6 AUG 02 CAPLUS and CA patent records enhanced with European and Japan
Patent Office Classifications
NEWS 7 AUG 02 The Analysis Edition of STN Express with Discover!
(Version 7.01 for Windows) now available
NEWS 8 AUG 04 Pricing for the Save Answers for SciFinder Wizard within
STN Express with Discover! will change September 1, 2004
NEWS 9 AUG 27 BIOCOMMERCE: Changes and enhancements to content coverage
NEWS 10 AUG 27 BIOTECHABS/BIOTECHDS: Two new display fields added for legal
status data from INPADOC
NEWS 11 SEP 01 INPADOC: New family current-awareness alert (SDI) available
NEWS 12 SEP 01 New pricing for the Save Answers for SciFinder Wizard within
STN Express with Discover!
NEWS 13 SEP 01 New display format, HITSTR, available in WPIDS/WPINDEX/WPIX
NEWS 14 SEP 14 STN Patent Forum to be held October 13, 2004, in Iselin, NJ
NEWS 15 SEP 27 STANDARDS will no longer be available on STN
NEWS 16 SEP 27 SWETSCAN will no longer be available on STN

NEWS EXPRESS JULY 30 CURRENT WINDOWS VERSION IS V7.01, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 11 AUGUST 2004
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 17:12:25 ON 27 SEP 2004

=> file ca,caplus,medline,biosis,wpids,drugu,uspatfull

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

1.26

1.26

FILE 'CA' ENTERED AT 17:16:09 ON 27 SEP 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'CAPLUS' ENTERED AT 17:16:09 ON 27 SEP 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'MEDLINE' ENTERED AT 17:16:09 ON 27 SEP 2004

FILE 'BIOSIS' ENTERED AT 17:16:09 ON 27 SEP 2004
Copyright (c) 2004 The Thomson Corporation.

FILE 'WPIDS' ENTERED AT 17:16:09 ON 27 SEP 2004
COPYRIGHT (C) 2004 THE THOMSON CORPORATION

FILE 'DRUGU' ENTERED AT 17:16:09 ON 27 SEP 2004
COPYRIGHT (C) 2004 THE THOMSON CORPORATION

FILE 'USPATFULL' ENTERED AT 17:16:09 ON 27 SEP 2004
CA INDEXING COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

=> s ophthalmic (5a) aceclidine
L1 18 OPTHALMIC (5A) ACECLIDINE

=> dup remove L1
PROCESSING COMPLETED FOR L1
L2 11 DUP REMOVE L1 (7 DUPLICATES REMOVED)

=> d L2 1-11 bib, ab

L2 ANSWER 1 OF 11 USPATFULL on STN
AN 2004:139468 USPATFULL
TI Composition for treatment of night sight problems(halos, comas and glare) after refractive surgery, intra ocular lens implant after lensectomy or intraocular implant in phakic patients comprising aceclidine employed at low concentrations
IN Randazzo, Alessandro, Milano, ITALY
PI US 2004106644 A1 20040603
AI US 2003-473740 A1 20031002 (10)
WO 2002-EP3542 20020329
PRAI IT 2001-MI708 20010403
DT Utility
FS APPLICATION
LREP SUGHRUE MION, PLLC, 2100 PENNSYLVANIA AVENUE, N.W., SUITE 800, WASHINGTON, DC, 20037
CLMN Number of Claims: 10
ECL Exemplary Claim: 1
DRWN 3 Drawing Page(s)
LN.CNT 342

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB After refractive surgery to reduce ametropia (i.e. myopia, astigmatism or hypermetropia) an average percentage of patients between 15.8% after PRK (Photo Refractive Keratectomy) and 33% after LASIK (Laser in situ Keratomileusis) shows a poor night sight due to the presence of halos, glare and coma. A comparable disorder is present in a percentage of patients that underwent lensectomy (cataract or refractive lensectomy) with intra ocular lens (IOL) implant and IOL implants in phakic patients

to reduce ametropia. Thanks to the effect on pupillary kinetics, diluted low concentrations (from 0.002% to 0.040%) of Aceclidine were surprisingly found to effectively reduce and/or eliminate night sight problems for about 6 hours.

L2 ANSWER 2 OF 11 CA COPYRIGHT 2004 ACS on STN DUPLICATE 1
 AN 137:273249 CA
 TI Composition for the treatment of night sight problems (halos, comas and glare) after refractive surgery, intra ocular lens implant after lensectomy or intraocular lens implant in phakic patients comprising aceclidine employed at low concentrations
 IN Randazzo, Alessandro
 PA Italy
 SO PCT Int. Appl., 19 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002080915	A2	20021017	WO 2002-EP3542	20020329
WO 2002080915	A3	20030103		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1377292	A2	20040107	EP 2002-727502	20020329
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
US 2004106644	A1	20040603	US 2003-473740	20031002
PRAI IT 2001-MI708	A	20010403		
WO 2002-EP3542	W	20020329		

AB Method is disclosed for the the treatment of night sight problems (halos, comas and glare) after ophthalmic surgery. After refractive surgery to reduce ametropia (i.e. myopia, astigmatism or hypermetropia) an average percentage of patients between 15.8% after PRK (Photo Refractive Keratectomy) and 33% after LASIK (Laser in situ Keratomileusis) shows a poor night sight due to the presence of halos, glare and coma. A comparable disorder is present in a percentage of patients that underwent lensectomy (cataract or refractive lensectomy) with intra ocular lens (IOL) implant and IOL implants in phakic patients to reduce ametropia. Thanks to the effect on pupillary kinetics, diluted low concns. (from 0.002% to 0.040%) of aceclidine were surprisingly found to effectively reduce and/or eliminate night sight problems for about 6 h.

L2 ANSWER 3 OF 11 CA COPYRIGHT 2004 ACS on STN DUPLICATE 2
 AN 136:123671 CA
 TI Ophthalmic formulation of a selective cyclooxygenase-2 inhibitory drug
 IN Kararli, Tugrul T.; Bandyopadhyay, Rebanta; Singh, Satish K.; Hawley, Leslie C.
 PA Pharmacia & Upjohn Company, USA
 SO PCT Int. Appl., 71 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 4

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE

PI WO 2002005815 A1 20020124 WO 2001-US22061 20010712
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,
UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,
BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

AU 2001075908 A5 20020130 AU 2001-75908 20010712
US 2002035264 A1 20020321 US 2001-904098 20010712
EP 1303271 A1 20030423 EP 2001-953462 20010712
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

JP 2004528267 T2 20040916 JP 2002-511747 20010712

PRAI US 2000-218101P P 20000713
US 2001-279285P P 20010328
US 2001-294838P P 20010531
US 2001-296388P P 20010606
WO 2001-US22061 W 20010712

OS MARPAT 136:123671

AB A pharmaceutical composition suitable for topical administration to an eye contains a selective COX-2 inhibitor or nanoparticles of a drug of low water solubility, at a concentration effective for the treatment and/or prophylaxis of a disorder in the eye, and 1 or more ophthalmically acceptable excipients that reduce rate of removal from the eye such that the composition has an effective residence time of 2-24 h. Also provided is a method of treating and/or preventing a disorder in an eye, the method comprising administering to the eye a composition of the invention. Thus, an ophthalmic nanoparticle suspension contained valdecoxib at 2.15 mg/g, 1.2% glycerin, 0.8% EDTA disodium salt, 4.0% Gelcarin GP-379NF, 0.21% SeaSpen PF and 0.82% Povidone.

RE.CNT 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 4 OF 11 CA COPYRIGHT 2004 ACS on STN DUPLICATE 3
AN 133:340234 CA
TI Ophthalmic compositions for the treatment of visual disorders characterized by a reduced contrast sensitivity
IN Boldrini, Enrico; Severino, Dario Ercole; Panelli, Giorgio; Bianchini, Pietro
PA Farmigea S.p.A., Italy
SO PCT Int. Appl., 28 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000064425	A2	20001102	WO 2000-IT151	20000414
	WO 2000064425	A3	20010412		
	W:				
	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ,				
	DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,				
	IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,				
	MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,				
	SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM,				
	AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE,				
	DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,				
	CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	EP 1173207	A2	20020123	EP 2000-927713	20000414
	EP 1173207	B1	20021106		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO

AT 227135 E 20021115 AT 2000-927713 20000414
PRAI IT 1999-RM259 A 19990426
WO 2000-IT151 W 20000414

AB Disclosed are ophthalmic compns. able to reduce the impairments of the visual function (like halos, glare and reduction of the night and twilight vision) resulting from keratectomy operations carried out both with laser and by conventional techniques, from intraocular lens implantation (cataract surgery) and also resulting from various chronic pathologies affecting the anterior segment of the eyeball, such as the degenerations of the corneal tissue, which compns. contain one or more miotic agents, such as cholinomimetic active agents and cholinesterase inhibitors, in combination with one or more hypertonic agents, such as sulfacetamide and derivs. thereof, sodium chloride, glucose and glycerol. The miotic agents are selected from the group consisting of pilocarpine, carbachol, acetylcholine, aceclidine, physostigmine, and salts thereof.

L2 ANSWER 5 OF 11 CA COPYRIGHT 2004 ACS on STN DUPLICATE 4
AN 129:103734 CA
TI Bioavailability of timolol and aceclidine after ocular instillation in the rabbit

AU Matera, M. G.; Lampa, E.; Imperatore, A.; Berrino, L.; Russo, F.; Boldrini, E.; Rossi, F.

CS Institute of Pharmacology and Toxicology - Faculty of Medicine and Surgery, Second University of Naples, Naples, 80138, Italy

SO Research Communications in Molecular Pathology and Pharmacology (1998), 100(1), 35-42

CODEN: RCMPE6; ISSN: 1078-0297

PB PJD Publications Ltd.

DT Journal

LA English

AB The bioavailability of timolol and aceclidine after the ocular instillation of each drug (timolol 0.5% or aceclidine 2%) or both combined (timolol 0.5% + aceclidine 2%) has been evaluated in rabbits. 15 Male albino rabbits were treated by the instillation of timolol and aceclidine alone or combined in the conjunctival sac of the right eye. Timolol concns. in humor aqueous were assayed at 10 min, 30 min, 1 h, 2 h, 4 h and 6 h after instillation by HPLC. Aceclidine was assayed by a pharmacodynamic method: pupillary diameter at the following time intervals 0 (basal value), 1 min, 5 min, 30 min, 1 h, 2 h, 4 h, 6 h after treatment. The results demonstrated that no differences in timolol and aceclidine bioavailability were found between simple-drug preps. and their combination.

RE.CNT 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 6 OF 11 BIOSIS COPYRIGHT (c) 2004 The Thomson Corporation. on STN

AN 1989:155577 BIOSIS

DN PREV198936077618; BR36:77618

TI GLAUCOMA TREATMENT IN 1988 DRUGS LASER OR SURGERY.

AU HAMARD H [Reprint author]

CS CENTRE HOSP NATL D'OPHTALMOL DES QUINZE-VINGTS, 28 RUE DE CHARENTON, 75571 PARIS, CEDEX 12

SO Semaine des Hopitaux, (1988) Vol. 64, No. 38-39, pp. 2522-2523.

Meeting Info.: CONTINUING EDUCATION COURSES OF BICHAT PITIE-SALPETRIERE, PARIS, FRANCE, SEPTEMBER 26-30, 1988. SEM HOP PARIS.

CODEN: SHPAAI. ISSN: 0037-1777.

DT Conference; (Meeting)

FS BR

LA FRENCH

ED Entered STN: 13 Mar 1989

Last Updated on STN: 13 Mar 1989

L2 ANSWER 7 OF 11 BIOSIS COPYRIGHT (c) 2004 The Thomson Corporation. on
STN
AN 1985:57387 BIOSIS
DN PREV198528057387; BR28:57387
TI MEDICAL TREATMENT CHANCES IN CLOSED ANGLE GLAUCOMA.
AU REIBALDI A [Reprint author]; CANTATORE F; GUERRIERO S
CS ISTITUTO CLINICA OCULISTICA, UNIV BARI
SO Bollettino di Oculistica, (1984) Vol. 63, No. 7-8, pp. 669-674.
ISSN: 0006-677X.
DT Article
FS BR
LA ITALIAN

L2 ANSWER 8 OF 11 BIOSIS COPYRIGHT (c) 2004 The Thomson Corporation. on
STN
AN 1983:178767 BIOSIS
DN PREV198375028767; BA75:28767
TI THE DUALISTIC EFFECT OF PILOCARPINE AT THE EYE.
AU PILZ J [Reprint author]; PILZ A
CS AUGENKLIN MED AKAD CARL GUSTAV CARUS, DDR-8019 DRESDEN, FETSCHERSTR 74
SO Folia Ophthalmologica (Leipzig), (1982) Vol. 7, No. 3, pp. 195-201.
CODEN: FOOPDZ. ISSN: 0323-4932.
DT Article
FS BA
LA GERMAN

AB The competitive dualism of pilocarpine observed on other animal muscles, which seems to oppose general clinical experiences, is confirmed on inner ocular muscles. Pilocarpine works as competitive antagonist of carbachol and aceclidine (glaucostat), when these occur in a concentration, the effect of which exceeds the intrinsic activity of pilocarpine. If the concentrations are below this value, pilocarpine works as a competitive agonist. The effect does not depend on the temporary consequence of application of the drugs. The main object of investigation was the isolated sphincter iridis of the cattle with an intrinsic activity of pilocarpine of 0.26, completed by experiments on the human sphincter and the ciliary muscle, in which the intrinsic activity of pilocarpine is higher.

L2 ANSWER 9 OF 11 BIOSIS COPYRIGHT (c) 2004 The Thomson Corporation. on
STN
AN 1978:154518 BIOSIS
DN PREV197865041518; BA65:41518
TI ULTRASONOGRAPHIC STUDY OF THE EFFECT OF DIFFERENT MIOTICS ON THE EYE COMPONENTS.
AU FRANCOIS J [Reprint author]; GOES F
CS CLIN OPHTALMOL, AKAD ZIEKENHUIS, DE PINTELAAN 135, B-9000 GENT, BELG
SO Ophthalmologica, (1977) Vol. 175, No. 6, pp. 328-338.
CODEN: OPHTAD. ISSN: 0030-3755.
DT Article
FS BA
LA ENGLISH

AB The effect of topical instillations of carbachol 3%, pilocarpine 2%, aceclidine 2%, aceclidine 2%-adrenaline [epinephrine] 1% and of the Ocusert delivery system was determined and compared in 151 eyes [human]. The depth of the anterior chamber, the thickness and the position of the lens, the length of the vitreous and the refraction were studied. Aceclidine has negligible side effects on the ocular components (maximal change of the anterior chamber 0.20 mm; maximal change of the lens thickness 0.14 mm; maximal myopisation: -1.5 D. Carbachol has the strongest side effects (maximal change of the anterior chamber 0.80 mm; maximal change of the lens thickness 0.80 mm; maximal myopisation -11.50D). Carbachol and pilocarpine may cause an important forward displacement of the lens with the risk of an angle-closure glaucoma in an eye with shallow anterior chamber.

L2 ANSWER 10 OF 11 CA COPYRIGHT 2004 ACS on STN DUPLICATE 5
 AN 81:99141 CA
 TI **Ophthalmic** bioavailability. I. Corneal penetration of
aceclidine (3-acetoxyquinuclidine) into the rabbit eye using a
 perfusion technique
 AU Brian, B.; Boltralik, J. J.; Thom, L.; Zeleznick, L. D.
 CS Sci. Technol. Div., Alcon Lab., Inc., Fort Worth, TX, USA
 SO Journal of Pharmaceutical Sciences (1974), 63(4), 633-5
 CODEN: JPMSAE; ISSN: 0022-3549
 DT Journal
 LA English
 AB A rapid, efficient, and sensitive method was developed for extraction and
 subsequent quantitation of aceclidine-HCl (3-acetoxyquinuclidine-HCl) (I)
 [6109-70-2] and 3-quinuclidinol-HCl [6238-13-7] from biol. fluid. After
 adjustment of pH and salt concentration, chloroform exts. of serum, urine, or
 aqueous
 fluid could be quantitated by gas-liquid chromatog. without derivative
 formation. The analytical procedure was used to determine the corneal
 absorption of I in the rabbit eye. I, a drug used topically in glaucoma
 treatment, entered the anterior chamber of the eye by penetration
 exclusively through the cornea. In expts. on conjunctival absorption, the
 amount of I found in the anterior chamber was <50 ng. The corneal
 absorption expts. gave cumulated absorption of 1-8.5 µg in 30 mins.

L2 ANSWER 11 OF 11 BIOSIS COPYRIGHT (c) 2004 The Thomson Corporation. on
 STN
 AN 1982:75331 BIOSIS
 DN PREV198223005323; BR23:5323
 TI OCULAR BIOMETRY.
 AU FRANCOIS J [Reprint author]
 CS DEP OPHTHALMOL, UNIV GHENT, DE PINTELAAN, 135, B-9000 GHENT, BELG
 SO Doc. Ophthalmol. Proc. Ser., pp. P135-164. THIJSSSEN, J. M. AND A. M.
 VERBEEK (ED.). DOCUMENTA OPHTHALMOLOGICA PROCEEDINGS SERIES, VOL. 29.
 ULTRASONOGRAPHY IN OPHTHALMOLOGY; PROCEEDINGS OF THE 8TH SIDUO (SOCIETAS
 INTERNATIONALIS PRO DIAGNOSTICA ULTRASONICA IN OPHTHALMOLOGIA) CONGRESS,
 NIJMEGEN, NETHERLANDS. XIV+538P. DR W. JUNK BV PUBLISHERS: THE HAGUE,
 NETHERLANDS. BOSTON, MASS., USA (DIST. IN THE USA BY KLUWER BOSTON, INC.:
 HINGHAM, MASS.). ILLUS. 1981 (RECD. 1982).
 Publisher: Series: Documenta Ophthalmologica Proceedings Series.
 CODEN: DOPSBP. ISSN: 0303-6405. ISBN: 90-6193-724-8.
 DT Book
 Conference; (Meeting)
 FS BR
 LA ENGLISH

=> s aceclidine and glare
 L3 11 ACECLIDINE AND GLARE

=> d 13 1-11 bib, ab

L3 ANSWER 1 OF 11 CA COPYRIGHT 2004 ACS on STN
 AN 137:273249 CA
 TI Composition for the treatment of night sight problems (halos, comas and
glare) after refractive surgery, intra ocular lens implant after
 lensectomy or intraocular lens implant in phakic patients comprising
aceclidine employed at low concentrations
 IN Randazzo, Alessandro
 PA Italy
 SO PCT Int. Appl., 19 pp.
 CODEN: PIXXD2
 DT Patent
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002080915	A2	20021017	WO 2002-EP3542	20020329
	WO 2002080915	A3	20030103		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	EP 1377292	A2	20040107	EP 2002-727502	20020329
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
	US 2004106644	A1	20040603	US 2003-473740	20031002
PRAI	IT 2001-MI708	A	20010403		
	WO 2002-EP3542	W	20020329		

AB Method is disclosed for the the treatment of night sight problems (halos, comas and glare) after ophthalmic surgery. After refractive surgery to reduce ametropia (i.e. myopia, astigmatism or hypermetropia) an average percentage of patients between 15.8% after PRK (Photo Refractive Keratectomy) and 33% after LASIK (Laser in situ Keratomileusis) shows a poor night sight due to the presence of halos, glare and coma. A comparable disorder is present in a percentage of patients that underwent lensectomy (cataract or refractive lensectomy) with intra ocular lens (IOL) implant and IOL implants in phakic patients to reduce ametropia. Thanks to the effect on pupillary kinetics, diluted low concns. (from 0.002% to 0.040%) of **aceclidine** were surprisingly found to effectively reduce and/or eliminate night sight problems for about 6 h.

L3 ANSWER 2 OF 11 CA COPYRIGHT 2004 ACS on STN
 AN 133:340234 CA
 TI Ophthalmic compositions for the treatment of visual disorders characterized by a reduced contrast sensitivity
 IN Boldrini, Enrico; Severino, Dario Ercole; Panelli, Giorgio; Bianchini, Pietro
 PA Farmigea S.p.A., Italy
 SO PCT Int. Appl., 28 pp.
 CODEN: PIXXD2
 DT Patent
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000064425	A2	20001102	WO 2000-IT151	20000414
	WO 2000064425	A3	20010412		
	W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
	EP 1173207	A2	20020123	EP 2000-927713	20000414
	EP 1173207	B1	20021106		
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
	AT 227135	E	20021115	AT 2000-927713	20000414

PRAI IT 1999-RM259 A 19990426
WO 2000-IT151 W 20000414

AB Disclosed are ophthalmic compns. able to reduce the impairments of the visual function (like halos, **glare** and reduction of the night and twilight vision) resulting from keratectomy operations carried out both with laser and by conventional techniques, from intraocular lens implantation (cataract surgery) and also resulting from various chronic pathologies affecting the anterior segment of the eyeball, such as the degenerations of the corneal tissue, which compns. contain one or more miotic agents, such as cholinomimetic active agents and cholinesterase inhibitors, in combination with one or more hypertonic agents, such as sulfacetamide and derivs. thereof, sodium chloride, glucose and glycerol. The miotic agents are selected from the group consisting of pilocarpine, carbachol, acetylcholine, **aceclidine**, physostigmine, and salts thereof.

L3 ANSWER 3 OF 11 CAPLUS COPYRIGHT 2004 ACS on STN

AN 2002:793416 CAPLUS

DN 137:273249

TI Composition for the treatment of night sight problems (halos, comas and **glare**) after refractive surgery, intra ocular lens implant after lensectomy or intraocular lens implant in phakic patients comprising **aceclidine** employed at low concentrations

IN Randazzo, Alessandro

PA Italy

SO PCT Int. Appl., 19 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002080915	A2	20021017	WO 2002-EP3542	20020329
	WO 2002080915	A3	20030103		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	EP 1377292	A2	20040107	EP 2002-727502	20020329
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
	US 2004106644	A1	20040603	US 2003-473740	20031002
PRAI	IT 2001-MI708	A	20010403		
	WO 2002-EP3542	W	20020329		

AB Method is disclosed for the the treatment of night sight problems (halos, comas and **glare**) after ophthalmic surgery. After refractive surgery to reduce ametropy (i.e. myopia, astigmatism or hypermetropia) an average percentage of patients between 15.8% after PRK (Photo Refractive Keratectomy) and 33% after LASIK (Laser in situ Keratomileusis) shows a poor night sight due to the presence of halos, **glare** and coma. A comparable disorder is present in a percentage of patients that underwent lensectomy (cataract or refractive lensectomy) with intra ocular lens (IOL) implant and IOL implants in phakic patients to reduce ametropy. Thanks to the effect on pupillary kinetics, diluted low concns. (from 0.002% to 0.040%) of **aceclidine** were surprisingly found to effectively reduce and/or eliminate night sight problems for about 6 h.

L3 ANSWER 4 OF 11 CAPLUS COPYRIGHT 2004 ACS on STN

AN 2000:772437 CAPLUS
 DN 133:340234
 TI Ophthalmic compositions for the treatment of visual disorders
 characterized by a reduced contrast sensitivity
 IN Boldrini, Enrico; Severino, Dario Ercole; Panelli, Giorgio; Bianchini,
 Pietro
 PA Farmigea S.p.A., Italy
 SO PCT Int. Appl., 28 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000064425	A2	20001102	WO 2000-IT151	20000414
	WO 2000064425	A3	20010412		
	W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
	EP 1173207	A2	20020123	EP 2000-927713	20000414
	EP 1173207	B1	20021106		
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
	AT 227135	E	20021115	AT 2000-927713	20000414
PRAI	IT 1999-RM259	A	19990426		
	WO 2000-IT151	W	20000414		

AB Disclosed are ophthalmic compns. able to reduce the impairments of the visual function (like halos, glare and reduction of the night and twilight vision) resulting from keratectomy operations carried out both with laser and by conventional techniques, from intraocular lens implantation (cataract surgery) and also resulting from various chronic pathologies affecting the anterior segment of the eyeball, such as the degenerations of the corneal tissue, which compns. contain one or more miotic agents, such as cholinomimetic active agents and cholinesterase inhibitors, in combination with one or more hypertonic agents, such as sulfacetamide and derivs. thereof, sodium chloride, glucose and glycerol. The miotic agents are selected from the group consisting of pilocarpine, carbachol, acetylcholine, aceclidine, physostigmine, and salts thereof.

L3 ANSWER 5 OF 11 WPIDS COPYRIGHT 2004 THE THOMSON CORP on STN
 AN 2002-740978 [80] WPIDS
 DNC C2002-209929
 TI Use of aceclidine in the treatment of night sight problems in patients who have undergone refractive surgery or intraocular phakic lens implant or intraocular implant in aphakic patients.
 DC B02
 IN RANDAZZO, A
 PA (RAND-I) RANDAZZO A
 CYC 101
 PI WO 2002080915 A2 20021017 (200280)* EN 19
 RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ NL OA PT SD SE SL SZ TR TZ UG ZM ZW
 W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK DM DZ EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ OM PH PL PT RO RU SD SE SG SI SK SL TJ TM TN TR TT TZ UA UG US UZ VN YU ZA ZM ZW

EP 1377292 A2 20040107 (200404) EN

R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT
RO SE SI TR

AU 2002257735 A1 20021021 (200433)

US 2004106644 A1 20040603 (200436)

ADT WO 2002080915 A2 WO 2002-EP3542 20020329; EP 1377292 A2 EP 2002-727502
20020329, WO 2002-EP3542 20020329; AU 2002257735 A1 AU 2002-257735
20020329; US 2004106644 A1 WO 2002-EP3542 20020329, US 2003-473740
20031002

FDT EP 1377292 A2 Based on WO 2002080915; AU 2002257735 A1 Based on WO
2002080915

PRAI IT 2001-MI708 20010403

AB WO 2002080915 A UPAB: 20021212

NOVELTY - Use of **aceclidine** (I) or one of its derivatives in the
manufacture of a composition for the treatment of night sight problems
(i.e. halos, coma and **glare**) in patients who have undergone
refractive surgery or intraocular phakic lens implant or intraocular
implant in aphakic patients, is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for:

(1) Pharmaceutical composition for the treatment of night sight
problems (i.e. halos, coma and **glare**) in patients who have
undergone refractive surgery or intraocular phakic lens implant or
intraocular implant in aphakic patients, comprising (I) and a carrier; and

(2) Method of treating night sight problems (i.e. halos, coma and
glare) in patients who have undergone refractive surgery or
intraocular phakic lens implant or intraocular implant in aphakic
patients, by topical administration of (I) in an ophthalmic composition.

ACTIVITY - Ophthalmic.

In a double-masked randomized clinical trial with 14 patients (27
eyes), it was found that 18 out of 19 eyes treated with **aceclidine**
ophthalmic compositions versus 2 out of 8 treated with placebo showed an
improvement in night vision problems (95 vs 25%). The efficacy was 6 hours
with onset after 15-20 minutes following instillation. Side effects were
modest and transient.

MECHANISM OF ACTION - None given.

USE - (I) is used in the treatment of night sight problems (i.e.
halos, coma and **glare**) in patients who have undergone refractive
surgery or intraocular phakic lens implant or intraocular implant in
aphakic patients.

ADVANTAGE - (I) solves the problem of light ray diffraction and
aberration during the night hours. Administration of an ophthalmic
composition containing (I) and in very low concentrations may effectively
reduce the pupillary diameter for a period of up to 6 hours. (I) has high
selectivity compared to other parasympathomimetic drugs, addressing the
effective reduction/prevention of halos, coma and glares in patients who
had refractive surgery, where no side effects were detectable.

Dwg.0/3

L3 ANSWER 6 OF 11 USPATFULL on STN

AN 2004:139468 USPATFULL

TI Composition for treatment of night sight problems(halos, comas and
glare) after refractive surgery, intra ocular lens implant after
lensectomy or intraocular implant in phakic patients comprising
aceclidine employed at low concentrations

IN Randazzo, Alessandro, Milano, ITALY

PI US 2004106644 A1 20040603

AI US 2003-473740 A1 20031002 (10)

WO 2002-EP3542 20020329

PRAI IT 2001-MI708 20010403

DT Utility

FS APPLICATION

LREP SUGHRUE MION, PLLC, 2100 PENNSYLVANIA AVENUE, N.W., SUITE 800,
WASHINGTON, DC, 20037

CLMN Number of Claims: 10

ECL Exemplary Claim: 1
DRWN 3 Drawing Page(s)
LN.CNT 342

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB After refractive surgery to reduce ametropia (i.e. myopia, astigmatism or hypermetropia) an average percentage of patients between 15.8% after PRK (Photo Refractive Keratectomy) and 33% after LASIK (Laser in situ Keratomileusis) shows a poor night sight due to the presence of halos, **glare** and coma. A comparable disorder is present in a percentage of patients that underwent lensectomy (cataract or refractive lensectomy) with intra ocular lens (IOL) implant and IOL implants in phakic patients to reduce ametropia. Thanks to the effect on pupillary kinetics, diluted low concentrations (from 0.002% to 0.040%) of **Aceclidine** were surprisingly found to effectively reduce and/or eliminate night sight problems for about 6 hours.

L3 ANSWER 7 OF 11 USPATFULL on STN

AN 2003:206917 USPATFULL

TI Medical uses of in situ formed gels

IN Viegas, Tacey X., Birmingham, AL, UNITED STATES

Reeve, Lorraine E., Dexter, MI, UNITED STATES

Henry, Raymond L., St. Clair Shores, MI, UNITED STATES

PI US 2003143274 A1 20030731

AI US 2002-234922 A1 20020904 (10)

RLI Continuation of Ser. No. US 2000-628227, filed on 28 Jul 2000, PENDING
Continuation of Ser. No. US 1999-330618, filed on 11 Jun 1999, GRANTED,
Pat. No. US 6136334 Continuation of Ser. No. US 1996-773755, filed on 23
Dec 1996, GRANTED, Pat. No. US 5958443 Continuation of Ser. No. US
1993-174101, filed on 28 Dec 1993, GRANTED, Pat. No. US 5587175 Division
of Ser. No. US 1991-785305, filed on 30 Oct 1991, GRANTED, Pat. No. US
5318780

DT Utility

FS APPLICATION

LREP Pillsbury Winthrop, LLP, Suite 200, 11682 El Camino Real, San Diego, CA,
92130

CLMN Number of Claims: 20

ECL Exemplary Claim: 1

DRWN No Drawings

LN.CNT 1147

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Balanced pH, hyperosmotic, hypoosmotic, or isoosmotic gels are ideal vehicles for drug delivery. They are especially suited for topical body cavity or injection application of drugs or diagnostic agents; for drug or diagnostic agent delivery to the eye of a mammal; as protective corneal shields; or as ablatable corneal masks useful in laser reprofiling of the cornea. The compositions without the addition of a drug or diagnostic agent are useful as medical devices, for instance, in separating surgically or otherwise injured tissue as a means of preventing adhesions.

L3 ANSWER 8 OF 11 USPATFULL on STN

AN 2000:141902 USPATFULL

TI Medical uses of in situ formed gels

IN Viegas, Tacey X., Birmingham, AL, United States

Reeve, Lorraine E., Dexter, MI, United States

Henry, Raymond L., St. Clair Shores, MI, United States

PA MDV Technologies, Inc., San Diego, CA, United States (U.S. corporation)

PI US 6136334 20001024

AI US 1999-330618 19990611 (9)

RLI Continuation of Ser. No. US 1996-773755, filed on 23 Dec 1996, now
patented, Pat. No. US 5958443 which is a continuation of Ser. No. US
1993-174101, filed on ~~28 Dec~~ 1993, now abandoned which is a continuation
of Ser. No. US 1991-785305, filed on 30 Oct 1991, now patented, Pat. No.
US 5318780

DT Utility
FS Granted
EXNAM Primary Examiner: Azpuru, Carlos
LREP Pillsbury Madison & Sutro, LLPW. Patrick BengtssonNan Wu
CLMN Number of Claims: 8
ECL Exemplary Claim: 1
DRWN No Drawings
LN.CNT 1137

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Balanced pH, hyperosmotic, hypoosmotic, or isoosmotic gels are ideal vehicles for drug delivery. They are especially suited for topical body cavity or injection application of drugs or diagnostic agents; for drug or diagnostic agent delivery to the eye of a mammal; as protective corneal shields; or as ablatable corneal masks useful in laser reprofiling of the cornea. The compositions without the addition of a drug or diagnostic agent are useful as medical devices, for instance, in separating surgically or otherwise injured tissue as a means of preventing adhesions.

L3 ANSWER 9 OF 11 USPATFULL on STN
AN 1999:117015 USPATFULL
TI Medical uses of in situ formed gels
IN Viegas, Tacey X., Canton, MI, United States
Reeve, Lorraine E., Dexter, MI, United States
Henry, Raymond L., Grosse Pointe Woods, MI, United States
PA MDV Technologies, Inc., San Diego, CA, United States (U.S. corporation)
PI US 5958443 19990928
AI US 1996-773755 19961223 (8)
RLI Continuation of Ser. No. US 1993-174101, filed on 28 Dec 1993, now patented, Pat. No. US 5587175 which is a continuation of Ser. No. US 1991-785305, filed on 30 Oct 1991, now patented, Pat. No. US 5318780

DT Utility
FS Granted
EXNAM Primary Examiner: Azpuru, Carlos A.
LREP Pillsbury Madison & Sutro LLP
CLMN Number of Claims: 39
ECL Exemplary Claim: 1
DRWN No Drawings
LN.CNT 1248

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Balanced pH, hyperosmotic, hypoosmotic, or isoosmotic gels are ideal vehicles for drug delivery. They are especially suited for topical body cavity or injection application of drugs or diagnostic agents; for drug or diagnostic agent delivery to the eye of a mammal; as protective corneal shields; or as ablatable corneal masks useful in laser reprofiling of the cornea. The compositions without the addition of a drug or diagnostic agent are useful as medical devices, for instance, in separating surgically or otherwise injured tissue as a means of preventing adhesions.

L3 ANSWER 10 OF 11 USPATFULL on STN
AN 96:118391 USPATFULL
TI Medical uses of in situ formed gels
IN Viegas, Tacey X., Canton, MI, United States
Reeve, Lorraine E., Dexter, MI, United States
Henry, Raymond L., Grosse Pointe Woods, MI, United States
PA MDV Technologies, Inc., Dearborn, MI, United States (U.S. corporation)
PI US 5587175 19961224
AI US 1993-174101 19931228 (8)
RLI Division of Ser. No. US 1991-785305, filed on 30 Oct 1991, now patented, Pat. No. US 5318780
DT Utility
FS Granted
EXNAM Primary Examiner: Azpuru, Carlos

LREP Banner & Witcoff, Ltd.
CLMN Number of Claims: 11
ECL Exemplary Claim: 1
DRWN No Drawings
LN.CNT 1104

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Balanced pH, hyperosmotic, hypoosmotic, or isoosmotic gels are ideal vehicles for drug delivery. They are especially suited for topical body cavity or injection application of drugs or diagnostic agents; for drug or diagnostic agent delivery to the eye of a mammal; as protective corneal shields; or as ablatable corneal masks useful in laser reprofiling of the cornea. The compositions without the addition of a drug or diagnostic agent are useful as medical devices, for instance, in separating surgically or otherwise injured tissue as a means of preventing adhesions.

L3 ANSWER 11 OF 11 USPATFULL on STN

AN 94:48963 USPATFULL

TI Medical uses of in situ formed gels

IN Viegas, Tacey X., Canton, MI, United States

Reeve, Lorraine E., Dexter, MI, United States

Henry, Raymond L., Grosse Pointe Woods, MI, United States

PA Mediventures Inc., Dearborn, MI, United States (U.S. corporation)

PI US 5318780 19940607

AI US 1991-785305 19911030 (7)

DCD 20081210

DT Utility

FS Granted

EXNAM Primary Examiner: Page, Thurman K.; Assistant Examiner: Azpuru, Carlos

LREP Dykema Gossett

CLMN Number of Claims: 9

ECL Exemplary Claim: 1

DRWN No Drawings

LN.CNT 1057

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB Balanced pH, hyperosmotic, hypoosmotic, or isoosmotic gels are ideal vehicles for drug delivery. They are especially suited for topical body cavity or injection application of drugs or diagnostic agents; for drug or diagnostic agent delivery to the eye of a mammal; as protective corneal shields; or as ablatable corneal masks useful in laser reprofiling of the cornea. The compositions without the addition of a drug or diagnostic agent are useful as medical devices, for instance, in separating surgically or otherwise injured tissue as a means of preventing adhesions.